

Screening of biocontrol agent by phytochemical, antifungal activity against yellow mold disease in *Arachis Hypogaea* L.

M Tharanya, R Sagaya Giri

Department of Botany, Kunthavai Nachiyar Govt. Arts College (W), Autonomous (Affiliated to Bharathidasan University Tiruchirappalli), Thanjavur, Tamil Nadu, India

Abstract

The threat of aflatoxin contamination in food commodities and its association with health risks in both animals and humans. In the present investigation seventeen microflora were isolated from different ground nut filed area as *Aspergillus flavus*, *A. terreus*, *A. niger*, *A. ocraceus*, *A. spinulosus*, *A. fumigatus*, *Trichoderma* sp., *Fusarium solani*, *Curvuloria geniculata*, *Curvuloria lunata*, *Fusarium moniliforme*, *Penicillium granulatum*, *Cephalosporium* sp., *F. oxysporium*, *Helminthosporium oryzae*, *Memnomilla* sp. and *Penicillium notatum*. They performed antagonistic activity of soil microflora against *Aspergillus flavus*. *Trichoderma viride* has more suppression results recorded compare other soil fungi. And also studied the antifungal activity of plant extracts against *A. flavus*. The highest zone of inhibition recorded in 100µl concentration of *Acalypha indica*, *Amaranthus viride*, *Boerhavia diffusa* and *Tephrosia purpurea* leaves methanolic extract. These plant extract act as a good source of antifungal activity.

Keywords: groundnut, fungal diversity, antagonistic activity, phytochemical analysis, antifungal activity

Introduction

Soil is the major component of earth ecosystem which comprises of organic matter, minerals, gases and large numbers of macro and microorganisms. The soil ecosystem is supported by several interactions among its physical, chemical and biological components (Buscot, 2005) [3]. Many biological processes take place in soil and determine functions that provide various services within ecosystems: turnover of organic matter, symbiotic and non-symbiotic atmospheric nitrogen fixation, denitrification, aggregation etc. (Chenu and Stotzky, 2002) [4]. Nachiyar. Rhizosphere is the narrow zone of soil surrounding the root where microbe populations are stimulated by root activities. Soil microflora plays a pivotal role in evaluation of soil conditions and in stimulating plant growth (Kiran Singh *et al.*, 1999) [10]. Microorganisms in the soil and rhizosphere are beneficial in increasing soil fertility and plant growth as they are involved in several biochemical transformations and mineralization activities in soils. Type of cultivation and crop management practices found to have greater influence on the activity of soil microflora. Fungi are an important component of soil microbiota more in abundance than bacteria, depending on soil depth and nutrient conditions. Different soils have specific fungal flora, but the majority of species found in them are cosmopolitan (Ainsworth and Sussman, 1968) [1]. Fungi are fundamental for soil ecosystem functioning (Warcup, 1951) [17], especially in forest and agricultural soils, they play a key role in many essential processes such as organic matter decomposition and elemental release by mineralization (Christensen *et al.*, 1989). It was estimated 1.5million fungal species are present in natural ecosystems, but only 5-10% has been described formally (Hawksworth, 2001). The aim of the present investigation is to isolate mycoflora from different crop

Fields, to study fungal diversity and to observe percentage contribution of different fungal species. The study involves isolation, identification and enumeration of fungal species from different crop fields in and around Nanjangud taluk of Mysore district. Since ages, the basic requirements of humans like food, shelter, clothing and medicine is dependent on plant diversity. Herbal medicine is the oldest form of healthcare known to human society (Barnes *et al.*, 2007) [2]. Plants produce a diverse array of bioactive molecules, making them rich sources of diverse type of medicine.

The secondary metabolites such as alkaloids, glycosides, flavonoids, volatile oils, etc are biosynthetically derived from primary metabolites as protective mechanisms against predation by many microorganisms, insects and herbivores or any pathogenic attack on plants. The antifungal activity of four plant extracts from (*Acalypha indica*), (*Amaranthus viride*), (*Boerhavia diffusa*) and (*Tephrosia purpurea*) was *In vitro* evaluated against *Aspergillus flavus* sp. The pathogenicity of *Aspergillus flavus* sp. on seedlings was confirmed using soil infestation technique under greenhouse conditions.

Materials and Methods

Diversity of soil fungi

Collection of soil samples

Rhizosphere soil samples were collected from different groundnut field of various Districts in Tamil Nadu (S1, S2, S3, S4, S5 and S6) Table-1. In each field 1kg of soil sample was collected from the surface area reaching about 10-15 cm depth and near the rhizosphere region of plants. Soils were collected in sterile polythene bags and sealed on the spot. Samples were stored in laboratory at 4°C until further analysis.

Table 1: Soil samples of three different districts

Sample	Districts	Places
S1	Pudukkottai	Narangiyapattu
S2		Regunathapuram
S3	Thanjavur	Nambivayal
S4		Perayurani
S5	Thiruvavarur	Edamalaiyur
S6		Vaduvur

Isolation of Mycoflora

Dilution plate technique described by Warcup (1951) [17] was used for the isolation of fungi from various rhizosphere soil samples. 10 grams of soil samples were suspended in 90 ml of distilled water. The flasks were shaken thoroughly in order to get uniform distribution of the soil particles. The soil suspensions were diluted in 10 fold increment from 10^{-3} to 10^{-5} . The Volume of 10 ml of soil sample suspension from each serial dilution was pipetted onto different melted, cooled culture media Potato Dextrose Agar (PDA) supplemented with 1% Streptomycin. The pH of the culture media was maintained at 5.5 being optimal for the growth and sporulation in a majority of fungi. Each culture media was prepared in a liter of distilled water and autoclaved at 120°C at 15 psi for 20 min. 1% Streptomycin was used as an antibiotic for the restrain of bacterial growth. Each colony was sub cultured and maintained on potato dextrose agar slants. The inoculated plates were incubated at room temperature $28 \pm 2^\circ\text{C}$ in an inverted position for 5-7 days. Three replicates were maintained for each sample.

Data Analysis

Population density is expressed in terms of Colony Forming Unit (CFU) per gram of soil with dilution factors. The percentage contribution of each isolate was calculated by using the following formula (Gaddeyya *et al.*, 2012) [6].

Identification of Soil Fungi

Fungal morphology were studied macroscopically by observing colony features (Texture and Color) and microscopically by staining with Lacto phenol cotton blue and observed under compound microscope for conidiophores, conidia and arrangement of spores. The fungi were identified with the help of literature (Nagamani *et al.*, 2006).

Effect of antagonistic fungi on the suppression of pathogenic fungi of yellow mold disease

The antagonistic activity of various soil fungi against the test pathogen were studied with dual culture plate technique (Morton and Strouble, 1955) [11] under *in vitro* condition. The test pathogen *Aspergillus flavus* and the soil fungi viz., *Aspergillus terreus*, *A. Niger*, *A. fumigatus*, *F. oxysporium*, *P. notatum*, *P. granulatatum* and *Trichoderma viride* were grown separately on PDA medium. The agar block cut from actively growing margin of individual species of soil fungi and test organism were inoculated opposite to each other approximately 3 cm apart on potato dextrose agar medium in petriplates. Three replicates were maintained for each set. Controls were set in single and dual inoculated cultures at the fungus. The position of the colony margin on the back of the disc was recorded daily. The assessment was made for both organisms. The percentage of inhibition of growth was calculated as follows.

$$\text{Percentage of inhibition of growth} = \frac{r - r'}{r} \times 100$$

Where,

r = growth of the fungus from the Centre of the colony towards the Centre of the plate in the absence of antagonistic fungus.

r' = growth of the fungus from the Centre of the colony towards the antagonistic fungus.

The colony interaction between test pathogen and soil fungi was proposed methods (Skidmore and Dickinson 1976) [15].

Qualitative analysis of phytochemical compounds

Phytochemical analysis of the all plant extracts was undertaken using standard qualitative methods as described by various authors. The plant extracts were screened for the presence of biologically active compounds such as alkaloids, flavonoids, carbohydrates, phytosterols, proteins, phenolics, tannins and saponins (Harborne, 1973).

Antifungal activity

Twenty grams of *Acalypha indica*, *Amaranthus viride*, *Boerhavia diffusa* and *Tephrosia purpurea* leaves were taken separately and surface sterilized with 70% alcohol and finally with sterilized distilled water. Then they were crushed by pestle and mortar and extracted with 20 ml of sterilized distilled water and filtered aseptically through double layered cheese cloth. The extracts were poured in the flasks plugged with cotton to avoid contamination. The extracts were diluted to different concentrations (25, 50, 75, 100%) by adding distilled sterile water for further use. Then obtained extract was subjected to antifungal assay by food poisoned technique (Nene and Thapliyal, 1993 and Grover and Moore, 1962) [12, 7].

Results and Discussion

Isolation and identification

The Present investigation was conducted to find out the fungal diversity in six different groundnut fields. In this study, 136 fungal colonies of 16 fungal species belonging to 7 genera were isolated and identified. Sixteen fungal isolates were identified (Table-2). *Aspergillus flavus*, *A. terreus*, *A. niger*, *A. ocraceus*, *A. spinulosus*, *A. fumigatus*, *Trichoderma* sp., *Fusarium solani*, *Curvularia geniculata*, *Penicillium granulatatum*, *F. oxysporium*, *Helminthosporium oryae*, *Memnomilla* sp. and *Penicillium notatum* based on cultural characters and sporulation structure. The colonies of *Aspergillus* and *Fusarium* belonging to were predominant in all soil samples in groundnut fields. The similar and significant report has already been made in all soil samples of crop fields such as sunflower, sesame, capsicum, rice, green gram, sugarcane, ground nut and black gram by Shiny Niharika *et al.*, (2013) [14].

Gaddeyya *et al.*, (2012) [6] determined the fungal population and their diversity in agricultural fields of Salur. During the investigation period 173 fungal colonies of 15 fungal species were observed. The maximum fungal species belongs to Deuteromycotina (169 colonies) and Zygomycotina (4 colonies) were observed. Among the isolates the genera *Aspergillus* and *Penicillium* were dominant. Rohilla and Salar (2012) [13] twenty three soil samples were characterized for the incidence of fungal strains from pesticides contaminated agricultural soils. A

total of 59 fungal isolates were obtained from the analyses of 23 soil samples taken from pesticide contaminated soils

through soil dilution agar plating, Warcup's soil plate and Waksman Direct inoculation method.

Table 2: Occurrence of soil mycoflora in different Groundnut field from various Districts of Tamilnadu

Sample	Pudukkottai		Thanjavur		Thiruvavur		Total No. of Colonies
	S1	S2	S3	S4	S5	S6	
<i>A. flavus</i>	2	1	3	2	4	2	14
<i>A. niger</i>	1	1	2	1	1	1	7
<i>A. terreus</i>	3	2	3	2	2	1	13
<i>A. ocraceus</i>	1	1	1	-	1	1	5
<i>A. spinulorsus</i>	1	-	-	2	-	3	6
<i>A. fumigatus</i>	1	1	2	2	1	2	9
<i>C. lunata</i>	1	-	4	-	-	-	8
<i>Cu. geniculata</i>	1	-	1	2	2	-	4
<i>Cepalosporium sp.</i>	2	2	1	2	3	-	12
<i>F. moniliforme</i>	2	2	2	2	-	3	12
<i>F. oxysporium</i>	2	2	1	1	2	2	10
<i>H. oryzae</i>	1	1	1	1	1	2	7
<i>Memnomilla sp</i>	1	1	1	1	-	1	5
<i>Penicillium sp</i>	1	1	1	-	1	1	5
<i>P. granulatam</i>	2	2	2	1	3	1	11
<i>Trichoderma sp</i>	2	2	1	2	1	1	8
Total No. of colonies	24	19	26	22	22	23	136

Effect of antagonistic fungi on the suppression of pathogenic fungi of yellow mold disease

In the present study, antagonistic activity of *Aspergillus terreus*, *A. Niger*, *A. fumigatus*, *F. oxysporium*, *P. notatum*, *P. granulatam* and *Trichoderma viride* against *Aspergillus flavus* were studied by *in vitro* dual culture experiment. The species of *Trichoderma* and *A. Niger* showed the ability to inhibit the pathogen. But these species showed variability in the percentage of inhibition. The percentage inhibition of growth of pathogen against *Trichoderma viride*, *Aspergillus Niger* and *A. fumigatus* were 45, 42 and 39 percentage with respectively. Growth inhibition decreased as follows; *Aspergillus terreus* (31) *A. Niger*, (42), *A. fumigatus*, (39) *F. oxysporium*, (34) *P. notatum*, (33) *P. granulatam*, (31) and *Trichoderma viride* (45) Table-3.

Swati *et al.*, (2018) [16] in dual culture *In vitro* antagonistic activity of *Trichoderma harzianum* was studied against soilborne fungal pathogens isolated from rhizosphere soil of groundnut viz. *Aspergillus flavus*, *Aspergillus niger*, *Fusarium roseum*, *Macrophomina phaseolina*, *Phythium myriotylum*, *Rhizoctonia solani* and *Sclerotium rolfsii*. *Trichoderma harzianum* had been marked as significant inhibition of selected fungal pathogens as compared to their respective controls. The different workers checked these pathogens against *Trichoderma harzianum* from different hosts. During the present investigation all these pathogens were firstly tested against *Trichoderma harzianum* isolated from rhizosphere soil of groundnut from Marathwada region.

The maximum percentage of inhibition were revealed by *Trichoderma harzianum* against *Fusarium roseum* (62.18%), followed by *Phythium* (51.85%), *Aspergillus flavus* (50.37%), *Aspergillus niger* (46.55%), *Rhizoctonia solani* (41.78%), *Macrophomina phaseolina* (30.58 %) and minimum inhibition was shown against *Sclerotium rolfsii* (27.73%).

Dhotre and Wanmare (2017) [5] were got the equal interaction of *Trichoderma* against root rot causing fungus of soybean i.e. *Fusarium moniliforme* (85 mm) followed by *Rhizoctonia solani* (81.5 mm), *Fusarium oxysporum* (80 mm), *Fusarium solani* (79 mm), *Fusarium roseum* (79 mm), *Phytophthora asojae* (76 mm), *Macrophomina phaseolina* (73.5 mm). Also the were found that the *Trichoderma harzianum* inhibit the growth of *Macrophomina phaseolina* followed by *Aspergillus Niger* and *Fusarium oxysporum* pathogens isolated from seeds. The *Trichoderma* reached the pathogen within 3- 4 days and overgrew them in 8-10 days. The occurrence of an inhibition zone in dual culture characterizes the secretion of some diffusible non-volatile substances. The zone of inhibition was found against two pathogens i.e. *Macrophomina phaseolina* and *Rhizoctonia solani*. This zone of inhibition lasts for 3 to 7 days and then this antagonist was overgrowing on them. Similar kind of results was found by Hajjigharari *et al.*, (2008) [8] for different species of *Trichoderma* including *Trichoderma harzianum* against *Rhizoctonia solani* and *Macrophomina phaseolina*.

Table 3: Effect of antagonistic fungi on the suppression of pathogenic fungi

S. No.	Growth response of the antagonistic and test fungus	Antagonistic fungi tested						
		<i>A. terreus</i>	<i>A. niger</i>	<i>A. fumigatus</i>	<i>F. oxysporium</i>	<i>P. notatum</i>	<i>P. granulatam</i>	<i>T. viride</i>
1	Colony growth of the pathogen towards antagonist(mm)	20	23	21	24	26	19	17
2	Colony growth of the pathogen away from the antagonist (mm)	19	22	21	21	23	22	24
3	% growth inhibition of the pathogen in the zone of the interaction (mm)	24	20	28	31	30	25	26
4	% Colony growth of the antagonist in control i.e. Growth towards the centre of plate in the absence of the pathogen (mm)	31	42	39	26	33	31	45
5	Control growth of the antagonist away from the pathogen (mm)	24	17	31	20	22	26	40
6	% growth of inhibition in the zone of interaction	20	22	19	34	24	27	20

Qualitative analysis of phytochemical compounds

Results obtained for qualitative screening of phytochemicals in the leaves of *Acalypha indica*, *Amaranthus viride*, *Boerhavia diffusa* and *Tephrosia purpurea* were presented in table-. Preliminary Phytochemical screening of the methanolic leaf extracts of *Acalypha indica*, reveals the presence of various biochemical compounds which include alkaloids, flavonoids, carbohydrates, proteins, phenols, saponins, tannins, triterpenoids, phytosterols and terpenoids. Phytochemical screening of the Methanol leaf extracts of

Amaranthus viride by qualitative study showed the presence of alkaloids, flavonoids, carbohydrates, proteins, saponins, tannins, triterpenoids, phytosterols and terpenoids. Methanol extracts of leaves of *Boerhavia diffusa* showed the presence of alkaloids, flavonoids, carbohydrates, proteins, tannins, terpenoids. Whereas phytosterols, triterpenoids, saponins and phenols were absent. Similarly *Tephrosia purpurea* showed the presence of alkaloids, flavonoids, carbohydrates, proteins, phenols, tannins, triterpenoids and terpenoids were absent. (Table-4)

Table 4: Qualitative analysis of the phytochemicals studies for the methanolic extract of selected plants

Phytochemical test	<i>Acalypha indica</i>	<i>Amaranthus viride</i>	<i>Boerhavia diffusa</i>	<i>Tephrosia purpurea</i>
Alkaloids	+	+	+	+
Flavonoids	+	+	+	+
Terpenoids	+	+	+	+
Triterpenoids	+	+	-	+
Phenols	+	-	-	+
Saponins	+	+	-	-
Protein	+	+	+	+
Tannin	+	+	+	+
Carbohydrate	+	+	+	+
Phytosterols	+	+	-	-

(+) – indicates presence, (-) – indicates absence

In vitro evaluation of plants extracts against *Aspergillus flavus*

Four leaves extract such as *Acalypha indica*, *Amaranthus viride*, *Boerhavia diffusa* and *Tephrosia purpurea* were significantly reduced the mycelia growth of *Aspergillus flavus* compared to control. All concentration of the leaf extract was found to be inhibitory to both growth and the rate of inhibition increased generally by increasing the concentration, 100% concentration was the most effective concentration. Table 5 showed the percentage of Radial growth of *Aspergillus flavus*. The growth of *Aspergillus flavus* was inhibited at 25% concentration of *Acalypha indica*, *Amaranthus viride*, *Boerhavia diffusa* and *Teprosia*

purpurea leaves extracts by 27.3±0.30, 24.6±0.24, 23.0±0.11 and 22.4±0.20 respectively. On the other hand, at 50% concentration of the foresaid plant extracts inhibited mycelial growth of *A. flavus* by 24.0±0.20, 23.4±0.22, 21.2±0.30 and 18.0±0.30 respectively where as in 75% concentration 21.8±0.31, 20.3±0.30, 18.6±0.20 and 16.3±0.21 of inhibition were recorded respectively. The highest concentration of 100% of *Acalypha indica*, *Amaranthus viride*, *Boerhavia diffusa* and *Tephrosia purpurea* leaf extracts showed effectiveness in minimizing the colony growth of fungi were 28.0±0.20, 26.0±0.23, 22.9±0.21 and 19.8±0.30 respectively.

Table 5: In vitro antifungal activity of four plant leaf in methanol extracts against *Aspergillus flavus*

Fungal organism	Concentration	Radial growth (mm)			
		<i>Acalypha indica</i>	<i>Amaranthus viride</i>	<i>Boerhavia diffusa</i>	<i>Tephrosia purpurea</i>
<i>A. flavus</i>	Control	73.3±0.21	73.3±0.21	73.3±0.21	73.3±0.21
	25	22.4±0.20	27.3±0.30	24.6±0.24	23.0±0.11
	50	18.0±0.30	23.4±0.22	21.2±0.30	24.0±0.20
	75	16.3±0.21	20.3±0.30	18.6±0.20	21.8±0.31
	100	19.8±0.30	22.9±0.21	26.0±0.23	28.0±0.20

Data represent the mean ± standard deviation of three replicates.

Conclusion

In this study, 136 fungal colonies of 16 fungal species belonging to 7 genera were isolated and identified from six different rhizosphere soil samples. *Aspergillus flavus* is the causative organisms of yellow mold disease. The fungus colonies are white in color and produce three types of spores, micro conidia, macro conidia and chlamydospores. In antagonistic activity, *Trichoderma viride*, *A.niger* had a marked significantly inhibitory effect on the growth of selected phytopathogens compared to their. The use of these bio-agents are not only safe for the farmers and consumers but also eco-friendly, cose effective, easy to produce and easy to apply the formulation. The plant extracts were screened for the presence of bio-compounds such as alkaloids, flavonoids, carbohydrates, proteins, phenols,

saponins, tannins, triterpenoids, phytosterols and terpenoids. There are four widely growing plants such as *Acalypha indica*, *Amaranthus viride*, *Boerhavia diffusa* and *Tephrosia purpurea* were used in the antifungal study by Food poison technique. All plant leaf extracts were found to be inhibiting the growth of the pathogen and the rate of inhibition increased generally by increasing the concentration. These studies form the base for eco-friendly management of plant disease which will minimize the usage of pesticide and chemical fungicide.

References

1. Anisworth GC, Sussman AS. The fungi as Advanced Treatise. The fungal Population. 1968; 3:426-496.

2. Barnes JW. Global-scale surface spectral variations on Titan seen from Cassini/VIMS, Icarus. 2007; 186:242-258, doi:10.1016/j.icarus.2006.08.021
3. Buscot F, Varma A. Micro-organisms in soils: Roles in genesis and functions. Soil Biology. Heidelberg. 2005; 3:3-17.
4. Chenu C, Stotzky G. Interactions between microorganisms and soil particles: an overview. In: Interactions between soil and particles and microorganisms, 2002, 205-218.
5. Dhotre Vanmare. Antagonistic activity of *Trichoderma* against root rot causing fungi of soybean. International Journal of Advanced Scientific Research. 2017; 2(2):57-58.
6. Gaddeyya G, Shiny Niharika P, Bharathi P, Ratna Kumar PK. Isolation and identification of soil mycoflora in different crop fields at Salur Mandal, Advances in Applied Science Research. 2012; 3(4):2020-2026.
7. Grover RK, Moore JD. Toximetric studies of fungicides against the brown rot organisms, *Sclerotiniafruticola* and *S. laxa*. Phytopathology. 1962; 52:876-880.
8. Hajighararari B, Torabi-giglou M, Mohammadi MR, Pavari M. Biological potential of some Iranian *Trichoderma* isolates in the control of soilborne plant pathogenic fungi. African Journal of Biotechnology. 2008; 7:967-972.
9. Hawksworth DL. Tropical Mycology. Micromycetes CABI. 2002; 2:1-11.
10. Kiran Singh, Jaishree Borana, Sobha Srivastava. Journal of soil biology and ecology. 1999; 19:11-14.
11. Morton DT, Stroube NH. Antagonistic and stimulatory effects of microorganism upon *Sclerotium rolfsii*. Phytopathology. 1955; 45:419-420.
12. Nene YL, Thapliyal PN. Fungicides in Plant Disease Control. Oxford and IBH Publishing Co., New Delhi. 1993; 45:552-579.
13. Rohilla SK, Salar RK. Isolation and characterization of various fungal strains from agricultural soil contaminated with pesticides. Research Journal of Recent Sciences. 2012; 1:297-303.
14. Shiny Niharika P, Gaddeyya G, Ratna Kumar PK. An investigation on soil mycoflora of different crop fields at Narasannapetamandal, Srikakulam district. International Research Journal of Environment Sciences. 2013; 2(9):38-44.
15. Skidmore AM, Dickinson CM. Colony interactions and hyphal interferences between *Septoria nodorum* and phylloplane fungi. Trans. Br. Mycol. Soc. 1976; 66:57-64.
16. Swati N Gaikwad, Sadhana N Salve, Suchita K Rajurkar. *In vitro* antagonistic activity of *Trichoderma harzianum* against soilborne fungal pathogens, International Journal of Biology Research. 2018; 3(2):87-89.
17. Warcup JH. The ecology of soil fungi, Trans Br Mycol Soc. 1951; 34:376-399.
18. Warcup JM. The Soil plate method for isolation of fungi from Soil, Nature, Lond, 1950, 117-166.